

The role of *Campsurus notatus* (Ephemeroptera: Polymitarcytidae) bioturbation and sediment quality on potential gas fluxes in a tropical lake

João José Fonseca Leal · André Luiz dos Santos Furtado ·
Francisco de Assis Esteves · Reinaldo Luiz Bozelli ·
Marcos Paulo Figueiredo-Barros

Published online: 7 April 2007
© Springer Science+Business Media B.V. 2007

Abstract About 30% of the total area of Lake Batata (Amazon) was impacted by the disposal of bauxite tailings originated from the process of washing bauxite. This effluent, composed by fine particles of clay and water, settled on top of the natural sediment, originating a new substratum with a different physical and chemical composition. This phenomenon created a new distinct habitat (impacted sediment) influencing the benthic community. The aim of this study was to evaluate the impact of bioturbation by *Campsurus notatus* (Ephemeroptera: Polymitarcytidae)

on potential gas fluxes in the sediment of natural and impacted areas of the lake. The natural sediment had a significantly higher methane concentration when compared to the impacted one. In incubated sediment cores, the presence of *C. notatus* nymphs resulted in a significant increase in oxygen consumption and methane and carbon dioxide release to the water column. The effect of the presence of nymphs on methane was ambiguous. The *C. notatus* nymphs strongly decreased methane concentration in natural sediment samples, probably because of the enhancement of the oxic sediment area. However, this effect was not observed in impacted samples. Finally, the new substratum of Lake Batata decreased methane concentration in sediment and water column. *C. notatus* nymphs demonstrated to have a significant role on gas flux (methane and CO₂) from sediment to water column as well as on oxygen consumption in Lake Batata, consequently influencing the carbon cycle in this lake.

Handling editor: S. M. Thomaz

J. J. F. Leal (✉)
Centro Federal de Educação Tecnológica de
Química – Nilópolis, Rua Lúcio Tavares, 1045,
Nilópolis, RJ 26530-060, Brazil
e-mail: joaoleal@biologia.ufrj.br

A. L. dos Santos Furtado
Empresa Brasileira de Pesquisa Agropecuária, Centro
Nacional de Pesquisa de Monitoramento por Satélite,
Av. Dr. Júlio Soares de Arruda, 803, Parque São
Quirino, Campinas, SP 13088-300, Brazil
e-mail: andre@cnpm.embrapa.br

F. de Assis Esteves · R. L. Bozelli ·
M. P. Figueiredo-Barros
Inst. de Biologia, Dep. de Ecologia, Laboratório de
Limnologia, Universidade Federal do Rio de Janeiro,
Cidade Universitária – Ilha do Fundão, CCS, Rio de
Janeiro 21940-590, Brazil

Keywords Amazonian lake · Bioturbation ·
Campsurus notatus · Methane · Sediment

Introduction

In the mid 70s the mining company Mineração Rio do Norte S.A. began its activity of bauxite ore

extraction at Porto Trombetas (Pará, Brazil). As part of the process, bauxite ore must be washed to remove clay particles. Until the end of 1989, bauxite tailings (water + clay) originated from the washing process was dumped directly into Lake Batata, impacting nearly 30% of the total area of the lake. This effluent is composed of solid particles (clay) and is deposited on the bottom of Lake Batata over its natural sediment formed by layers of up to 5 m. The new substratum is poor in organic matter, nutrients, and has extremely fine granulometry ($<0.49 \mu\text{m}$).

By the end of the 80s several studies were carried out in this aquatic ecosystem searching for solutions that could mitigate and understand the impact of this disturbance (eg. Bozelli et al., 2000). The deposition of clay particles in Lake Batata altered the physical and chemical aspects of the sediment (Roland & Esteves, 1993; Callisto & Esteves, 1996), leading to a decrease in the richness and abundance of many benthic species (Leal & Esteves, 1999).

Nymphs of *Campsurus notatus* (Needham and Murphey 1924 Ephemeroptera: Polymitarcyidae) have colonized the new habitat at the impacted area. The occurrence of this organism in higher densities and biomass in the impacted area, as compared to the natural one, indicates that this species has benefited from the conditions imposed by the presence of bauxite tailings (Leal & Esteves, 1999; 2000).

The nymphs build long U-type tunnels in sediment where a continuous water flow is established, revealing a high bioturbation capacity in Amazonian floodplain lakes, as well as at other tropical aquatic ecosystems (Cressa, 1986; Nolte, 1987; Melo et al., 1993). The importance of *C. notatus* to bacterioplankton production was recently demonstrated because of its effect on the mineralization and liberation of nitrogen stocked in the sediment to the water column (Leal et al., 2003). Nevertheless, until now, the effect of these organisms on carbon cycling and organic matter decomposition is unclear.

The aim of this research was to experimentally evaluate the effect of *C. notatus* nymph colonization on aerobic and anaerobic organic matter decomposition (methanogenesis) in both natural and impacted sediment. This will provide more

direct and conclusive information about nutrient cycling and the role of benthic fauna (post disturbance) on Lake Batata sediment.

Study area

Located in the Amazonian plain, on the right margin of Trombetas River, Lake Batata is situated between $1^{\circ}25'$ and $1^{\circ}35'$ S and $56^{\circ}15'$ and $56^{\circ}25'$ W in Porto Trombetas, District of Oriximiná (State of Pará, Brazil). Lake Batata is classified as a clear water ecosystem (Sioli, 1984), with high transparency, low electric conductivity, low nutrient concentration, predominantly slightly acid or acid pH (Bozelli et al., 2000). Some limnological and morphological data were described by Ferrão-Filho & Esteves (1994) and Panosso et al. (1995), respectively. Biological features were investigated by Bozelli (1992, 1994) who researched the zooplankton community. Anésio et al. (1997) and Huszar & Reynolds (1997) studied the bacterioplankton and phytoplankton communities, respectively.

This ecosystem shows high water level fluctuation associated to the Trombetas River flood pulse, causing a high variation in its depth (around 9.0 m) and surface area (from 18 Km^2 to 30 Km^2 approximately), representing low and high water periods (Panosso et al., 1995).

During 11 years (1979–1989), about 50,000 $\text{m}^3 \text{d}^{-1}$ of the resulting effluent originated from the bauxite washing process were dumped into Lake Batata. The liquid effluent is composed of solid particles (7%), predominantly clay, rich in silicate (47%), iron oxide and aluminum (21%), which was deposited on top of the natural sediment of the lake, covering approximately 3 km^2 of its total area. Tables 1 and 2 show a summarized description of the sediment granulometry and porosity, as well as nutrients and water content on natural and impacted areas.

Methods

Sediment sampling and acclimatization

During the low water period (December 2001), 0.03 m^3 of sediment from the 0–20 cm fraction

Table 1 Comparative sediment granulometric composition and porosity of the superficial sediment (0–10 cm; $n = 5$) from impacted and natural areas of Lake Batata

Sediment	Grain size (μm)	Impacted Sediment (%)	Natural Sediment (%)
Fine sand	62–125	0.93	1.35
Large silt	31–62	0.0	0.35
Medium silt	16–31	6.65	0.35
Fine silt	8–16	7.67	7.9
Very fine silt	4–8	8.85	4.5
Large clay	2–4	5.45	18.55
Medium clay	1–2	8.9	35.6
Fine clay	0.5–1	61.55	31.4
Porosity (%)		70.51	89.94

Table 2 Organic matter, total nitrogen, total carbon, and pore water concentration in the sediment of Lake Batata (Pará, Brazil)*

	Organic matter (%DW)	Total nitrogen (%DW)	Total carbon (%DW)	Pore water (%)
Natural area	20.80	0.46	5.10	73.86
Impacted area	14.80	0.18	1.72	62.03

* modified from Leal et al. (2003)

was sampled in both natural and impacted areas with a core sampler ($\varnothing = 8$ cm; area = 50 cm²) modified from Ambühl & Bühler (1975).

In the laboratory, natural and impacted sediment were washed in a 0.5 mm mesh size sieve to remove macroinvertebrates which were stored in glass flasks. Filtered lake water (GF/C Whatman) was added to the sediment, which was kept in the dark for 10 days at a mean temperature of 29.8°C (± 0.5).

The experiment consisted of 3 steps: acclimatization, pre-incubation and incubation. After acclimatization subsamples were taken to determine methane concentration and pore water content in sediment from natural and impacted areas (Fig. 1).

Sediment pre-incubation

Pre-incubation consisted of storing 20 cores from the acclimatized natural sediment and 20 from the acclimatized impacted sediment in thermal boxes. Each core ($L = 20$ cm; $\varnothing = 4$ cm) was filled with

8–10 cm of water and 10–12 cm of sediment and sealed in the bottom. All 40 samples were kept in filtered water from each sampling site. Aeration pumps were used to keep O_2 saturation near 100% (99.6 ± 0.5) both for inner and outer water. Water column was homogenized according to the model proposed by Svensson & Leonardson (1996) (Fig. 2).

During pre incubation (24 h) the cores were kept in the dark. After this period, 5 cores of each area were taken to determine dissolved oxygen concentration, CO_2 and CH_4 in water (initial concentration, see flux/consumption equation) and CH_4 in the sediment (0–2; 2–4; 4–6; 6–8 and 8–10 cm), and the pore water content.

Sediment incubation with *C. notatus* nymphs

The other 30 cores were divided into 2 groups (15 with natural sediment; 15 with impacted sediment), and were later divided into 3 subgroups of 5 cores each. Sub group 1 represented the control (without *C. notatus* nymphs). One or two nymphs were added in cores of subgroups 2 and 3 respectively, representing the density registered at Lake Batata (793 and 1587 ind. m^{-2}). After the introduction of the nymphs, the top of the cores were kept open for 3 h, and then, sealed for 6 h (incubation period). The CH_4 , CO_2 , and O_2 concentrations in the water and the CH_4 vertical distribution in the sediment were estimated at the end of the incubation period.

CH_4 and CO_2 sediment-water flux and O_2 consumption in the water were evaluated by the equation described by Dalsgaard et al. (2000).

Sediment – water flux/consumption

$$= ([F_c - I_c]) * V/a * t$$

where F_c = Final concentration; I_c = Initial concentration; V = Water volume (L); a = Core area (m^2) and t = incubation time (h).

Analyses

Dissolved oxygen concentration was estimated following the method proposed by Roland et al. (1999). CO_2 concentration was determined by the pH-alkalinity method (Carmouze, 1994).




Fig. 1 Diagram of the experiment represents pre-incubation phase and incubation phase. During pre-incubation the sediment samples from natural and impacted areas were saturated with O_2 for 6 h and incubated for 24 h in

the dark at $30^\circ C$. After pre-incubation the cores were kept open for 3 h. Incubation was stopped 6 h after atmosphere isolation and introduction of *Campsurus notatus*

To determine methane concentration in the pore water, 5 ml ($n = 5$) of each sediment fraction was taken and put in 12 ml flasks with 2 ml of NaOH (4%) and immediately sealed with rubber covers. Headspace subsamples were used to determinate methane concentration, which were kept in dark and low temperature conditions ($<10^\circ C$) until injection. Methane was determined by gas chromatography using a Varian Star 3400 chromatograph equipped with a POROPAK-Q column (1 m, 60/100 mesh) at $85^\circ C$, FID detector at $220^\circ C$ and injection at $120^\circ C$, N_2 was the carrier gas. The samples were injected using a Valvo C6W6 port loop valve (250 μ l). For data recording and peak registration a Star Chroma-

tography workstation 5.51 (Varian, USA) was used.

For the determination of water content and porosity, subsamples of fresh sediment were weighed in ceramic vessels and their weight loss recorded after heating for 4 days at $60^\circ C$ (Dalsgaard et al., 2000).

A two-way ANOVA Statistical analysis was performed on the acclimatized and pre-incubated sediment samples in order to assess the influence of depth and sediment type on methane concentration and to test the influence of *C. notatus* density (control, 1-nymph and 2-nymphs) on methane concentration. Differences between the sediment-water flux on controls and treatments




Fig. 2 Schematic representation of (a) pre-incubation phase (without *Campsurus notatus* and (b) incubation phase (with *Campsurus notatus*)

(1-nymph or 2-nymphs) were tested with the Kruskal-Wallis test with a Dunn's post test.

Results

Acclimatization, pre-incubation and incubation

Methane concentration profiles estimated from sediment after acclimatization and pre-incubation are shown in Fig. 3. We found a similar pattern on both natural and impacted sampling areas, i.e., the concentration increased significantly with depth (Two-way ANOVA, $F = 36.2$, $p < 0.05$). The effect of sediment type was strongly pronounced in the upper layers (0–2 cm) of natural and impacted sediment. The methane concentration in the natural top layer was 7–8 times higher than in the impacted one, contrasting with the

other layers, in which we observed a relatively lower difference (approximately 3 times less), but statistically significant (Two-way ANOVA, $F = 146.01$, $p < 0.05$).

Methane concentration of impacted sediment varied from $14.0 \mu\text{mol l}^{-1}$ to $110.8 \mu\text{mol l}^{-1}$ and from $10.0 \mu\text{mol l}^{-1}$ to $112.0 \mu\text{mol l}^{-1}$ after acclimatization and pre-incubation, respectively (Fig. 3a). The two-way ANOVA indicated no significant differences in methane concentration between the two treatments (Two-way ANOVA, $F = 0.25$, $p < 0.05$). On the other hand, we observed a statistically significant difference in the natural sediment (Two-way ANOVA, $F = 7.89$, $p < 0.05$), demonstrating that the artificial water oxygenation column between acclimatization and pre-incubation promoted a decrease of the methane concentration in the sediment (Fig. 3b). Nonetheless, the concentrations measured during acclimatization and pre-incubation




Fig. 3 Methane profile in sediment samples without nymphs of *Campsurus notatus*. Impacted sediment (a) and natural sediment (b). ● Acclimatization; □ Pre-incubation. Bars represent standard deviations ($n = 5$)

were in the range of amplitude usually observed in situ, both in natural and impacted sediments ($50\text{--}300 \mu\text{mol l}^{-1}$).

In the experiment with impacted sediment, the presence of *C. notatus* nymphs did not have significant effect on methane concentration (Two-way ANOVA, $F = 3.55$, $p < 0.05$; Fig. 4a), but methane concentration increased with depth. We recorded methane concentrations varying from $13.3 \mu\text{mol l}^{-1}$ to $119.9 \mu\text{mol l}^{-1}$ in controls; from $10.2 \mu\text{mol l}^{-1}$ to $116.7 \mu\text{mol l}^{-1}$ in cores incubated with 1 nymph; and from $16.0 \mu\text{mol l}^{-1}$ to $139.3 \mu\text{mol l}^{-1}$ in cores incubated with 2 nymphs.

Moreover, the results revealed a significant decrease in methane concentration in the natural cores with the addition of *C. notatus* nymphs (Two-way ANOVA, $F = 27.4$, $p < 0.05$; Fig. 4b). Methane concentration in control cores was significantly higher than those in the treatments with 1 or 2 nymphs. However, the statistical analysis showed no significant difference on methane concentration in natural sediment colonized by 1 or 2 nymphs (Two-way ANOVA, $F = 1.22$, $p < 0.05$)

Methane and carbon dioxide flux and oxygen consumption.

A significant increase in methane sediment-water flux was observed for both types of sediment (natural and impacted), independently of the density of nymphs, when compared to controls (Kruskal-Wallis test, $KW = 28.1$, $p < 0.05$; Table 3). For instance, in control samples of natural sediment the estimated flux was low ($0.52 \mu\text{mol m}^{-2} \text{h}^{-1}$), while in the core with 1 nymph the flux was 307 times higher, and in the cores with 2 nymphs it was 457 times higher. Methane flux in the 2-nymphs sample only was 1.5 times higher than in the 1-nymph sample.

CO₂ sediment-water flux was significantly enhanced in samples submitted to bioturbation when compared to controls for both types of sediment (Kruskal-Wallis test, $KW = 25.32$, $p < 0.05$; Table 3). CO₂ flux with 1 nymph, in treatments with either natural or impacted sediments, was 8 times higher than measured in the controls. In treatments with 2 nymphs, a significant increase in CO₂ flux was also observed. In




Fig. 4 Methane profile in sediment samples during the incubation with nymphs of *Campsurus notatus*. **(a)** Impacted sediment and **(b)** natural sediment. Nymphs

were kept in the cores for 6 h in the dark at 30°C. □ Control; ▲ 1 individual; △ 2 individuals. Bars represent standard deviations ($n = 5$)

contrast with the methane flux, the CO₂ sediment-water flux was not significantly different when comparing impacted and natural samples (Kruskal-Wallis test, $KW = 25,32$, $p < 0.05$). In natural sediment samples the flux varied from 0.11 (control) to 1.12 mmol m⁻² h⁻¹ (2-nymphs sample), while in impacted sediment samples the flux fluctuated from 0.09 (control) to 1.06 mmol m⁻² h⁻¹ (2-nymphs sample), indicating a proportional increase in CO₂.

Oxygen consumption displayed a similar pattern to the one observed for CO₂ flux, being significantly lower in the controls (absolute values) than in samples with one or two nymphs, independently of the type of sediment (Kruskal-Wallis test, $KW = 24.01$, $p < 0.05$; Table 3). Otherwise, in both natural and impacted sediments we did not observe a significant difference between treatments with one or two nymphs (Kruskal-Wallis test, $KW = 24.01$, $p < 0.05$). Oxygen consumption and carbon dioxide flux were less affected by the presence of nymphs than methane flux, as demonstrated by flux ratio and

sample ratio. Nymphs induced a raise in oxygen consumption and CO₂ flux, on average 3 and 12 times greater, respectively, whereas sample ratio measured for methane was higher.

Discussion

The development of research in Lake Batata was only made possible because of the unusual anthropogenic impact that occurred in this lake—the effluent of bauxite washing, which impacted a large area of the lake (nearly 30%). As a result of the dumping of the effluent, which ceased in 1989, part of the “igapó” (flooded forest) became “terra-firme” (nonflooded forest), causing death of the natural vegetation. On the other hand, some areas of the littoral zone, which in the past were inundated only at high water periods, are nowadays permanent flooded areas (Esteves, 2000).

Because of the colloidal and unconsolidated features of the clay, there are still bauxite tailings

Table 3 Methane and carbon dioxide flux, oxygen consumption in the water column and flux ratio (natural/impacted) and sample ratio (sample/control) in samples incubated with sediment from impacted and natural areas of Lake Batata (Pará, Brazil). Control represents samples without the presence of *C. notatus*. Samples were incubated in the dark at 30°C for 6 h (mean \pm standard deviation are shown; $n = 5$)

CH ₄ ($\mu\text{mol m}^{-2} \text{ h}$) CO ₂ (mmol m ² h) O ₂ (mmol m ² h)	Area	Control		1 Nymph		2 Nymphs		Sample ratio
		Control	Flux ratio	1 Nymph	Flux ratio	2 Nymphs	Flux ratio	
Impacted ¹ Natural	Impacted ¹ Natural	0.31 (0.16)	1.7	67.88 (30.75)	2.4	129.20 (48.21)	1.8	219–417
		0.52 (0.58)	1.2	159.60 (3.92)	1.3	237.84 (36.39)	1.1	307–457
Natural	Impacted	0.09 (0.01)	1.2	0.69 (0.11)	1.3	1.06 (0.23)	1.1	8–12
		0.11 (0.03)	1.3	0.87 (0.11)	1.5	1.12 (0.09)	1.6	8–10
Impacted	Natural	0.46 (0.12)	1.3	1.46 (0.18)	1.5	1.50 (0.21)	1.6	3–3
		0.60 (0.15)	1.3	2.17 (0.30)	1.5	2.43 (0.36)	1.6	4–4

suspended in the water column, with huge consequences for plankton (Bozelli, 1994; Anésio et al., 1997; Huszar & Reynolds, 1997) and fish communities (Lin & Caramaschi, 2005). However, the sediment was the most affected compartment since the disturbance destroyed some ecological niches and created others. In this new situation, *Campsurus notatus* nymphs have shown to be more capable of colonizing the sediment than other species of the benthic local fauna.

Data related to the effects of bauxite tailings on microorganisms are scarce and are basically restricted to bacterioplankton (Anésio, 2000; Farjalla et al., 2002) and nitrogen fixation and denitrification in sediment (Esteves et al., 2001). Furthermore, the scientific information concerning the genus *Campsurus* is restricted to seasonal dynamics and biomass (Takeda & Grzybkowska, 1997; Leal & Esteves, 2000) and due to the nature of the disturbance on Lake Batata there is no comparative data.

The lower methane concentrations in sediment with bauxite tailings indicate the influence of the sediment constitution on CH₄ production. The anthropogenic impact caused a decrease in nutrient availability such as C, N, and P (Roland et al., 2000), as well as changes in the physical structure of the sediment (Callisto, 2000) and water content. In general, the concentration of carbon in the impacted area is at least 7 times lower than that observed in the natural area.

Methane concentration depends on substratum availability of low molecular weight molecules such as acetate, methanol, formate, propanol and hydrogen (Angenent et al., 2002; Tonouchi, 2002; Weijma et al., 2003; Bodegom et al., 2004; Vetó et al., 2004), methane oxidation (Segers, 1998), and oxygen diffusion since this gas has a inhibitory capacity on methanogens (Fetzer et al., 1993; Frenzel et al., 1999) and it is also important as a substrate for aerobic oxidation. At Lake Batata, despite the fact that both areas present similar oxygen concentrations (about 0.25 mmol l⁻¹) in their superficial sediment layers (0–0.4 mm), the impact caused huge differences on the oxygen vertical profile. At the impacted area, oxygen penetrates twice as much as in the natural area, reaching 6 mm in depth, reflecting the fast oxygen consumption in the surface of natural sediment as

compared to the non organic impacted sediment (Leal et al., 2005). This fact might explain why methane concentration in the upper natural sediment layers was 7–8 times higher than in the impacted area. However, in the anoxic layers of the sediment, the natural/impacted concentration ratio was 3:4. Besides, methane vertical profile concentrations for both sediments were similar, i.e., the concentration gradually increases from the surface to the deepest layers of the sediment, which is an outcome usually found in continental aquatic environments (Mathias, 2000; Casper et al., 2003; Koizumi et al., 2003).

Benthic invertebrates, which build galleries or tubes, interfere in microbiological processes, feed bacteria and create structures with different properties from those observed in adjacent sediments, since they pump the water-sediment layer rich in oxygen through these tubes (Aller, 2001). The oxygen-rich water pumped by macrofauna, from top to bottom sediment layers, influences metabolic processes dynamics because it changes the oxi-reduction sites (Altmann et al., 2004) and stimulates the colonization of these galleries by aerobic bacteria and meiofauna (Wang et al., 2001). Kajan & Frenzel (1999) reported that Chironomidae larvae created microhabitats by constructing U-type or J-type tubes and increasing the bacterial activity in rice cultivation in Italy. Similarly, *C. notatus* nymphs built long galleries (more than 10 cm long), where oxygen concentration varied from 0.2 mmol l^{-1} to 0.3 mmol l^{-1} , the same oxygen concentration which was measured in the superficial sediment layer (Leal et al., 2005). This pattern had a strong effect on methane oxidation in the natural sediment of Lake Batata (Fig. 4b). Furthermore, our results indicate the importance of *C. notatus* nymphs in decreasing the methane concentration in sediments, probably both through induction of oxidation and release of methane into the water column. Nevertheless, the absence of perceptible methane bubbles in the sediment suggests that oxidation is the principal process involved. Recently, Kiyashko et al. (2004) demonstrated that *Stictochironomus pictulus* (Diptera: Chironomidae) larvae feed preferentially on methane-oxidizing bacteria, yet why this species has this selective feeding remains unclear.

In contrast, nymphs had no effect on methane concentration in impacted sediment (Fig. 4a). Part of this result might be explained by sediment granulometry. Clay deposition has changed mainly the physical structure, increasing the percentage of fine clay in the impacted sediment and changing its porosity, which might represent an impediment for gas diffusion (Table 1). Indeed, there is evidence that methane oxidation is positively related with soil particle size, likely complicating the diffusion of gas in soils with small granules (Min et al., 2002).

The influence of bioturbation observed only in the sediment from the natural area (Fig. 4b) contrasts with the flux results (Table 3), since we measured a raise of gas flux and gas consumption in both natural and impacted sediments. Bioturbation caused an increase of methane concentrations in the water column, which must have been due to the pumping of pore water with high CH_4 concentration. Svensson (1997) showed that the presence of *Chironomus plumosus* larvae stimulated the ammonium flux from the sediment to the overlying water, in part due to the pumping process. In addition, some authors reported that benthic fauna stimulated microbial activity in marine (Hansen & Kristensen, 1997; Heilskov & Holmer, 2001) and freshwater sediments (Svensson, 1997). Thus, it seems acceptable to infer that in Lake Batata *C. notatus* furthers methane production in sediment and increases methane concentration in the water column as a consequence of this pumping.

Oxygen consumption and carbon dioxide flux were less affected by the presence of nymphs than methane flux, as demonstrated by flux ratio and sample ratio, indicating that the activity of *C. notatus* nymphs stimulated a more effective anaerobic decomposition of organic carbon, even though it increases the oxygenation of the sediment during pumping. The sample ratio for methane varied from 219 to 457, while for the other gases the variation was lower (from 3 to 12). It is still unclear why methane flux was more affected by this benthic organism than carbon dioxide flux or oxygen consumption.

We assume that our results demonstrate the actual conditions in Lake Batata because both in situ and in the experimental study the

estimated methane concentration in situ were similar (data not shown) and *C. notatus* displayed the same behavior.

Even though our study describes a local phenomenon, it provides important information for other ecosystems because bauxite tailings are essentially composed by clay and clay is an important component of Amazonian and other tropical soils. Many other types of anthropogenic impacts may have similar effects by increasing sediment inflow. In a smaller scale but in the long term, farming and deforestation can have a similar outcome.

Finally, this study reveals that the new substratum (impacted sediment) of Lake Batata, which resulted from the deposition of bauxite tailings on top of the natural sediment layer, affected methane dynamics, leading to a lower methane concentration in the sediment and water column, and had a stronger effect on anaerobic organic matter mineralization.

Bauxite tailing influenced carbon cycle in two distinct ways: on one hand *C. notatus* nymphs contributed to a significant increase of CH₄ flux from the sediment to the water column and, on the other, stimulated methane oxidation in the sediment. In contrast, its influence on the organic matter aerobic mineralization was less clear. The results reveal a significant effect of *C. notatus* density on the fluxes of CH₄ and CO₂ as well as on O₂ consumption in both areas.

The bioirrigated zone produced by *C. notatus* raises microbiological activity in both types of sediment. This is proved by the high flux of CO₂ from the sediment to the water column. The results indicate that bioturbation promotes the colonization of the bioirrigated area of the impacted sediment by aerobic bacteria and intensifies the microbiological activity in this area. This implies that this species plays a role as a direct and indirect source of carbon incorporation into the sediment of lake thus improving ecological conditions in the impacted area of Lake Batata.

Acknowledgements The authors would like to thank Mineração Rio do Norte for the financial support and its staff for assisting in the sampling. We would also like to thank Dr. Peter Casper for his suggestions.

References

- Aller, R. C., 2001. Transport and reactions in the bioirrigated zone. In Boudreau, B. P. & B. B. Jørgensen (eds), *The Benthic Boundary Layer: Transport Processes and Biogeochemistry*. Oxford University Press, USA, 269–301.
- Altmann, D., P. Stief, R. Amann & D. de Beer, 2004. Nitrification in freshwater sediments as influenced by insect larvae: quantification by microsensors and fluorescence in Situ hybridization. *Microbial Ecology* 48: 145–153.
- Ambühl, H. & H. Bühner, 1975. Technik der Entnahme ungestörter Grossproben von Seesedimenten: ein verbessertes Bohrlot. *Schweizer Zeitschrift für Hydrologie* 37: 175–186.
- Anésio, A. M., 2000. Bacterioplankton. In Bozelli, R. L., F. A. Esteves & F. Roland (eds), *Lago Batata: impacto e Recuperação de um Ecossistema Amazônico*. IB-UFRJ/SBL, Rio de Janeiro, 73–88.
- Anésio, A. M., P. C. Abreu & F. A. Esteves, 1997. Influence of the hydrological cycle on the bacterioplankton of an impacted clear water Amazonian lake. *Microbial Ecology* 34: 66–73.
- Angenent, L. T., S. Sung & L. Raskin, 2002. Methanogenic population dynamics during startup of a full-scale anaerobic sequencing batch reactor treating swine waste. *Water Research* 36: 4648–4654.
- Bodegom, P. M., J. C. M. Scholten & A. J. M. Stams, 2004. Direct inhibition of methanogenesis by ferric iron. *FEMS Microbiology Ecology* 49: 261–268.
- Bozelli, R. L., F. A. Esteves & F. Roland (eds), 2000. *Lago Batata: Impacto e Recuperação de um Ecossistema Amazônico*. IB/UFRJ, Rio de Janeiro.
- Bozelli, R. L., 1994. Zooplankton community density in relation to water level fluctuations and inorganic turbidity in a Amazonian lake, Lago Batata, State of Pará, Brazil. *Amazoniana* 13: 17–32.
- Bozelli, R. L., 1992. Composition of the zooplankton community of Batata and Mussurá lakes and of the Trombetas river, State of Pará, Brazil. *Amazoniana* 12: 239–261.
- Callisto, M. & F. A. Esteves, 1996. Macroinvertebrados bentônicos em dois lagos amazônicos: lago Batata (um ecossistema impactado por rejeito de bauxita) e lago Mussurá (Brasil). *Acta Limnologica Brasiliensia* 8: 137–147.
- Callisto, M., 2000. Macroinvertebrados bentônicos. In Bozelli, R. L., F. A. Esteves & F. Roland (eds), *Lago Batata: impacto e Recuperação de um Ecossistema Amazônico*. IB/UFRJ, Rio de Janeiro: 139–151.
- Carmouze, J. P., 1994 *O metabolismo dos ecossistemas aquáticos: fundamentos teóricos, métodos de estudo e análises químicas*. Ed. Edgard Blücher/FAPESP, São Paulo, SP, Brasil.
- Casper, P., O. C. Chan, A. L. dos S. Furtado & D. D. Adams, 2003. Methane in an acidic bog lake: the influence of peat in the catchment on the biogeochemistry of methane. *Aquatic Sciences* 65: 36–46.

- Cressa, C., 1986. Variación estacional, distribución especial y balance energético de *Campsurus* sp. (Ephemeroptera: Polymitarcyidae) em el lago de Valencia, Venezuela. *Acta Científica Venezolana* 37: 572–579.
- Dalsgaard, T. (ed.), L. P. Nielsen, V. Brotas, P. Viaroli, G. Underwood, D. B. Nedwell, K. Sundbäck, S. Rysgaard, A. Milles, M. Bartoli, L. Dong, D. C. O. Thornton, L. D. M. Ottosen, G. Castaldelli & N. Risgaard-Petersen, 2000. Protocol handbook for NICE—Nitrogen Cycling in Estuaries: a project under the EU research programme: Marine Science and Technology (MAST III). National Environmental Research Institut, Silkeborg, Denmark.
- Esteves, F. A., 2000. Princípios Ecológicos para Mitigação do Impacto Antrópico. In Bozelli, R. L., F. A. Esteves, & F. Roland (eds), Lago Batata: Impacto e Recuperação de um Ecossistema Amazônico. IB-UFRJ/SBL, Rio de Janeiro, 1–16.
- Esteves, F. A., A. Enrich-Prast & D. D. Biesboer, 2001. Potential denitrification in submerged natural and impacted sediments of Lake Batata, an Amazonian lake. *Hydrobiologia* 444: 111–117.
- Farjalla, V. F., F. A. Esteves, R. L. Bozelli & F. Roland, 2002. Nutrient limitation of bacterial production in clear water Amazonian ecosystems. *Hydrobiologia* 489: 197–205.
- Ferrão-Filho, A. & F. A. Esteves, 1994. Nutritive value and sedimentation rates of particulate matter in the course of two flood pulses in na Amazonian Várzea lake. *Archiv für Hydrobiologie* 130: 325–337.
- Fetzer, S., F. Bak & R. Conrad, 1993. Sensitivity of methanogenic bacteria to oxygen and desiccation from paddy soil. *FEMS Microbiology Ecology* 12: 107–115.
- Frenzel, P., U. Bosse & P. H. Janssen, 1999. Rice roots and methanogenesis in a paddy soil: ferric iron as an alternative electron acceptor in the rooted soil. *Soil Biology and Biochemistry* 31: 421–430.
- Hansen, K. & E. Kristensen, 1997. Impact of macrofaunal recolonization on benthic metabolism and nutrient fluxes in a shallow marine sediment previously overgrown with macroalgal mats. *Estuarine Coastal and Shelf Science* 45: 613–628.
- Heilskov, A. C. & M. Holmer, 2001. Effects of benthic fauna on organic matter mineralization in fish-farm sediments: importance of size and abundance. *ICES Journal of Marine Science* 58: 427–434.
- Huszar, V. L. M. & C. S. Reynolds, 1997. Phytoplankton periodicity and sequences of dominance in an Amazonian flood-plain lake (Lago Batata, Pará, Brasil): responses to gradual environmental change. *Hydrobiologia* 346: 169–181.
- Kajan, R. & P. Frenzel, 1999. The effect of chironomid larvae on production, oxidation and fluxes of methane in a flooded rice soil. *FEMS Microbiology Ecology* 28: 121–129.
- Kiyashko, S. I., A. B. Imbs, T. Narita, V. I. Svetashev & E. Wada, 2004. Fatty acid composition of aquatic insect larvae *Stictochironomus pictulus* (Diptera: Chironomidae): evidence of feeding upon methanotrophic bacteria. *Comparative Biochemistry and Physiology (Part B)* 139: 705–711.
- Koizumi, Y., S. Takii, M. Nishino & T. Nakajima, 2003. Vertical distributions of sulfate-reducing bacteria and methane-producing archaea quantified by oligonucleotide probe hybridization in the profundal sediment of a mesotrophic lake. *FEMS Microbiology Ecology* 44: 101–108.
- Leal, J. J. F. & F. A. Esteves, 1999. Density and biomass of *Campsurus* sp. (Ephemeroptera) and other macroinvertebrates in an Amazonian lake impacted by bauxite tailings (Lago Batata, Pará, Brazil). *Amazoniana* 15: 193–209.
- Leal, J. J. F. & F. A. Esteves, 2000. Life cycle and production of *Campsurus notatus* (Ephemeroptera, Polymitarcyidae) in an Amazonian lake impacted by bauxite tailings (Pará, Brazil). *Hydrobiologia* 437: 91–99.
- Leal, J. J. F., F. A. Esteves, V. F. Farjalla & A. Enrich-Prast, 2003. Effect of *Campsurus notatus* on NH₄, DOC fluxes, O₂ uptake and bacterioplankton production in experimental microcosms with sediment-water interface of an Amazonian lake impacted by bauxite tailing. *International Review of Hydrobiology* 88: 167–178.
- Leal, J. J. F., A. Enrich-Prast, F. A. Esteves, R. Bozelli & V. F. Farjalla, 2005. Influence of *Campsurus notatus* bioturbation on oxygen profile and uptake in sediments of an Amazonian lake impacted by bauxite tailings. *Archiv für Hydrobiologie* 162(4): 557–574.
- Lin, D. S. C. & E. P. Caramaschi, 2005. Responses of the fish community to the flood pulse and siltation in a floodplain lake of the Trombetas River, Brazil. *Hydrobiologia* 75–91.
- Mathias, K., 2000. Methane turnover in exposed sediments of an Amazon floodplain lake. *Biogeochemistry* 50: 195–206.
- Melo, S. M., A. M. Takeda & N. C. Büttow, 1993. Variação temporal de ninfas de *Campsurus vialaceus* Needham and Murphy 1924 (Ephemeroptera: Polymitarcyidae) do Rio Baía (MS, Brasil). *Unimar (supp.)* 15: 95–107.
- Min, H., Z. Y. Chen, W. X. Wu & M. C. Chen, 2002. Microbial aerobic oxidation of methane in paddy soil. *Nutrient Cycling in Agroecosystems* 64: 79–85.
- Nolte, U., 1987. *Campsurus notatus* (Polymitarcyidae, Ephemeroptera) a bioturbation in Várzea lakes. *Amazoniana* 10: 219–222.
- Panosso, R. F., D. Muehe & F. A. Esteves, 1995. Morphological characteristics of na Amazon flood-plain lake (Lake Batata, Pará State, Brazil). *Amazoniana* 13: 245–258.
- Roland, F. & F. A. Esteves, 1993. Dynamics of phosphorus, carbon and nitrogen in Amazonian lake impacted by bauxite tailing (Batata Lake, Pará, Brazil). *Proceedings of the International Association of Theoretical and Applied Limnology* 25: 925–930.
- Roland, F., N. Caraco, J. J. Cole & P. del Giorgio, 1999. Rapid and precise determination of dissolved oxygen by spectrophotometry: evaluation of interference from color and turbidity. *Limnology and Oceanography* 44: 1148–1154.

- Roland, F., C. M. L. Ferreira & L. Kubrusly, 2000. Nutrientes e composição química do sedimento. In Bozelli R. L., F. A. Esteves & F. Roland (eds), Lago Batata: Impacto e Recuperação de um Ecossistema Amazônico. IB-UFRJ/SBL, Rio de Janeiro, 205–215.
- Segers, R., 1998. Methane production and methane consumption: a review of processes underlying wetland methane fluxes. *Biogeochemistry* 41: 23–51.
- Sioli, H., 1984. The Amazon and its main affluents: Hydrography, morphology of the river courses, and river types. In Sioli, H. (ed), *The Amazon: limnology and landscape ecology of a mighty tropical river and its basin*. Ed. Junk Publishers, Dordrecht, 127–165.
- Svensson, J. M., 1997. Influence of *Chironomus plumosus* larvae on ammonium flux and denitrification (measured by the acetylene blockage- and the isotope pairing-technique) in eutrophic lake sediment. *Hydrobiologia* 346: 157–168.
- Svensson, J. M. & L. Leonardson, 1996. Effects of bioturbation by tube-dwelling chironomid larvae on oxygen uptake and denitrification in eutrophic lake sediments. *Freshwater Biology* 35: 289–300.
- Takeda, A. M. & M. Grzybkowska, 1997. Seasonal dynamics and production of *Campsurus violaceus* nymphs (Ephemeroptera, Polymitarcyidae) in the Baía River, upper Paraná River floodplain, Brazil. *Hydrobiologia* 356: 149–155.
- Tonouchi, A., 2002. Isolation and characterization of a motile hydrogenotrophic methanogen from rice paddy field soil in Japan. *FEMS Microbiology Letters* 208: 239–243.
- Vető, I., I. Futó, I. Horváth & Z. Szántó, 2004. Late and deep fermentative methanogenesis as reflected in the H–C–O–S isotopy of the methane-water system in deep aquifers of the Pannonian Basin (SE Hungary). *Organic Geochemistry* 35: 713–723.
- Wang, F., A. Tessier & L. Hare, 2001. Oxygen measurements in the burrows of freshwater insects. *Freshwater Biology* 46: 317–327.
- Weijma, J., T. M. Chi, L. W. H. Pol, A. J. M. Stams & G. Lettinga, 2003. The effect of sulphate on methanol conversion in mesophilic upflow anaerobic sludge bed reactors. *Process Biochemistry* 38: 1259–1266.