

Effects of aluminum, calcium and low pH on egg hatching and nymphal survival of *Cloeon triangulifer* McDunnough (Ephemeroptera: Baetidae)*

Lisa M. Tabak¹ & K. Elizabeth Gibbs*

Department of Entomology, University of Maine, Orono, ME 04469, USA; ¹present address: Department of Biology, Queen's University, Kingston, Ontario, K7L 3N6, Canada (*author for correspondence)

Received 14 March 1990; in revised form 31 July 1990; accepted 10 October 1990

Key words: acidification; Ephemeroptera; pH; Al; Ca

Abstract

Laboratory experiments were conducted to assess the effects of aluminum, calcium and low pH on egg hatching and nymphal survival of the mayfly *Cloeon triangulifer*. Percent successful hatch (living nymphs breaking free of the chorion) decreased and percent partial hatch (nymphs dying attached to the chorion) increased with increasing acidity (pH 7.5–3.0). Most hatches occurring below pH 5.0 were partial hatches. Decreased time of exposure to acidic waters increased percent successful hatch and decreased percent partial hatch. Time to first hatch was not affected by pH. Eggs were incubated in acidic waters (pH 4.0 and 5.5) with additions of calcium (10 and 100 mg l⁻¹) and aluminum (100 and 500 µg l⁻¹). Aluminum decreased percent successful hatch and increased percent partial hatch and calcium increased both percent successful hatch and percent partial hatch. Time to first hatch was increased by both aluminum and calcium. The 96 h LC50 for small nymphs was pH 4.75. Addition of aluminum (100 and 500 µg l⁻¹) to acidic waters (pH 4.0 and 5.0) reduced nymphal mortality by 8–22%. Toxic effects of low pH on egg hatching and early nymphs may contribute to the absence of mayflies from acidified habitats.

Introduction

Mayflies are often reduced in abundance or absent in aquatic habitats at low pH (Weiderholm, 1984; Okland & Okland, 1986), although species such as *Leptophlebia vespertina* are tolerant of acidic conditions as low as pH 4.0. Decreased recruitment of *Ephemerella funeralis* after experimental stream acidification to pH 4.0

suggests that either eggs or small nymphs may be especially sensitive (Fiance, 1978). Successful egg hatching of *Leptophlebia cupida*, *Habrophlebia vibrans* and *Stenonema femoratum* has been recorded at pH 4.0 but significant mortality of nymphs of *Baetis flavistriga* occurred before the hatch was complete at pH 4.0 and 4.5 (Rowe *et al.*, 1988b). Effects on aquatic organisms by elevated hydrogen ion concentration may be modified by levels of aluminum (Baker & Schofield, 1982; Ormerod *et al.*, 1987; McCahon & Pascoe, 1989) and calcium (Brown, 1983). In addition to long-term low pH conditions, acid

*Contribution No. 1469 of the Maine Agricultural Experiment Station, University of Maine, Orono, Maine 04469 USA.

snow melt and severe storm events may produce temporary depressions in pH (Keller, 1983); therefore, the ability of an organism to survive may depend on tolerance of a particular life stage or the duration to which it is exposed.

Our objectives were to assess, with laboratory bioassays, the sensitivity of mayfly eggs and young nymphs to low pH and to determine whether aluminum, calcium, or both, negatively or positively modified the effect of low pH. Measures of effects on eggs were hatching rates and duration of embryonic development. Lethality experiments were conducted with young nymphs. The parthenogenic mayfly *Cloeon triangulifer* was selected for our experiments because previous experience indicated eggs having a high rate of hatching success could readily be obtained and nymphs could be successfully maintained in the laboratory (Gibbs, 1977).

Materials and methods

Chemistry of test waters

Water used in bioassays (dilution water) was double-distilled, deionized water titred with dechlorinated tap water to a specific conductance of 15–17 $\mu\text{S cm}^{-1}$, typical of surface waters in New England (Haines & Akielaszek, 1983). To avoid the use of chemical buffers, stock solutions were used to change the water in each bioassay container daily. Levels of pH were monitored daily, and adjustments were made in stock solutions only, using NaOH or H_2SO_4 . Aluminum was added as $\text{Al}_2(\text{SO}_4)_3$ and calcium as $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$; accordingly, mean sulfate concentrations increased with Al concentrations and mean chloride concentrations increased with Ca concentrations (Table 1).

Inflection point alkalinity was calculated using the Gran procedure (Stumm & Morgan, 1981). For cation determination, a 60 ml aliquot of water was acidified with 1.5 ml 1:1 HNO_3 and analyzed with a Perkin-Elmer model 703 atomic absorption spectrophotometer. Sodium and potassium were analyzed by air-acetylene flame,

calcium and magnesium by nitrous oxide-acetylene flame, and aluminum (total acid-reactive aluminum) by graphite furnace. Aluminum was also speciated into labile and non-labile monomeric fractions by passing a 60 ml aliquot through a cation exchange column (Dowex 50W-X8, Na form; procedure modified from Rosseland & Skogheim, 1982) and subsequently analysed as above. Anion (chloride, sulfate, nitrate, and fluoride) samples were filtered (0.45 μm Nucleopore) and frozen until analysed with a Dionex System 2110i ion chromatograph.

Egg bioassays

Eggs were obtained from adults reared from field-collected nymphs. Nymphs were collected from a pond with a pH of 7.5 ($\text{Ca}^{2+} = 13.2 \text{ mg l}^{-1}$; $\text{Mg}^{2+} = 4.4 \text{ mg l}^{-1}$; $\text{Cl} = 2.8 \text{ mg l}^{-1}$; $\text{SO}_4^{2-} = 6.5 \text{ mg l}^{-1}$; total Al = $49 \mu\text{g l}^{-1}$; specific conductance = $710 \mu\text{g l}^{-1}$) and transported to the laboratory in chilled containers where they were maintained in aerated glass dishes at 21 °C with a 16 h light : 8 h dark photoperiod. Leaf litter and associated organic sediment from the collection pond were provided as substrate and nourishment. Mature nymphs were placed under emergence cages. Subimagoes emerged from late afternoon through early evening, molted to the imaginal form in about 24 h, and were ready to oviposit 24 h later. Eggs were obtained by holding the mayfly by the wings with forceps and touching the tip of the abdomen to the surface of water in a petri dish; 150–1500 eggs were extruded by each mayfly. Eggs produced filamentous attachments to the substrate within 24 h of oviposition.

Eggs were incubated in test waters under static conditions in 60 × 15 mm petri dishes. A minimum of 100 eggs were used in each replicate and a minimum of three replicates were used in each experiment. Successfully hatched, partially hatched and dead eggs were counted and removed daily. In a successful hatch, a living nymph broke completely free from the chorion. In a partial hatch, the chorion was split to some degree but the nymph died without freeing itself. Eggs in

Table 1. Composition of dilution water adjusted for pH, aluminum, and calcium.

pH	Nominal Aluminum ($\mu\text{g l}^{-1}$)	Nominal Calcium (mg l^{-1})	Na ⁺ (mg l^{-1})	K ⁺ (mg l^{-1})	Ca ²⁺ (mg l^{-1})	Mg ²⁺ (mg l^{-1})	Total Al ($\mu\text{g l}^{-1}$)	Non-labile Al ($\mu\text{g l}^{-1}$)	Labile Al ($\mu\text{g l}^{-1}$)	F ⁻ (mg l^{-1})	Cl ⁻ (mg l^{-1})	NO ₃ ⁻ (mg l^{-1})	SO ₄ ²⁻ (mg l^{-1})
4.0	0	0	0.28	0.04	1.13	0.06	6	0	6	0.23	1.95	0	4.27
4.5	0	0	1.11	0.08	1.14	0.06	1	0	1	0.25	2.02	0	8.65
5.0	0	0	1.14	0.07	1.07	0.06	0	0	0	0.25	2.02	0	2.79
5.5	0	0	1.26	0.03	1.09	0.06	0	0	0	0.21	1.84	0	8.50
4.0	100	0	0.82	0.03	1.09	0.06	137	70	67	0.08	2.02	0	5.19
4.5	100	0	1.02	0.04	1.08	0.06	138	74	64	0.23	1.88	0	10.18
5.0	100	0	1.42	0.03	1.12	0.06	164	55	109	0.23	1.88	0	9.41
5.5	100	0	2.42	0.03	1.11	0.06	121	53	68	0.23	1.91	0	4.76
4.0	500	0	1.17	0.07	1.00	0.08	547	264	283	0.21	1.84	0	9.61
4.5	500	0	1.40	0.08	1.09	0.07	568	262	306	0.21	1.88	0	9.22
5.0	500	0	1.58	0.06	1.09	0.06	551	297	254	0.21	1.84	0	10.47
5.5	500	0	2.26	0.07	1.15	0.06	576	206	370	0.19	1.88	0	7.97
4.0	0	10	0.86	0.12	12	0.09	8	0	8	0.21	21.10	0	3.51
4.0	0	100	1.31	0.09	106	0.35	4	0	4	0.25	180.43	0	4.08
5.5	0	10	0.80	0.05	14	0.08	0	0	0	0.21	24.65	0	1.54
5.5	0	100	2.60	0.08	100	0.32	0	0	0	0.17	178.83	0	2.11
4.0	100	10	1.81	0.11	12	0.06	142	69	73	0.15	19.50	0	6.53
4.0	100	100	1.94	0.08	103	0.34	89	60	29	0.25	193.12	0	5.38
5.5	100	10	1.67	0.05	12	0.07	128	66	62	0.21	20.85	0	3.46
5.5	100	100	4.64	0.14	101	0.33	83	60	23	0.19	181.52	0	4.61
4.0	500	10	1.33	0.06	14	0.07	604	242	362	0.17	25.43	0	8.21
4.0	500	100	2.32	0.07	103	0.34	444	230	214	0.10	181.21	0	9.99
5.5	500	10	2.51	0.05	14	0.08	385	90	295	0.17	25.43	0	5.14
5.5	500	100	2.59	0.26	99	0.32	277	44	233	0.15	181.84	0	5.09

which no developing embryo could be detected and the egg content appeared vacuolated were considered dead. 'Days to first hatch' was used as a measure of the duration of embryonic development (Sweeney & Vannote, 1984).

Experiment A

To determine the effect of pH on egg hatching and duration of embryonic development, newly oviposited eggs were immediately exposed to test waters of pH 3.0, 4.0, 5.0, and 6.0. Controls consisted of dilution water adjusted to pH 7.5, that of the pond water from which the mayflies were collected.

Experiment B

The effects of delayed exposure to low pH were determined with a second group of eggs incubated at pH 7.5 for 7 d or until reaching the 'eyed' stage, indicating that hatching was imminent. The eggs were then exposed to the same pH levels as in Experiment A.

Experiment C

To examine effects of aluminum on egg hatching within the critical pH range determined in Experiment A, freshly oviposited eggs were incubated at pH 4.0, 4.5, 5.0, and 5.5 with Al concentrations of 0 (control), 100, and 500 $\mu\text{g l}^{-1}$.

Experiment D

Possible modifying effects of aluminum and calcium were examined by immediately exposing eggs at pH 4.0 and 5.5 with Al concentrations of 0 (control), 100, and 500 $\mu\text{g l}^{-1}$ and Ca concentrations of 0 (control), 10, and 100 mg l^{-1} .

Nymphal bioassays

Small (1–3 mm) field-collected nymphs were acclimated to dilution water for 24 h without food. Aerated, static 96 h bioassays were conducted in 300 cc glass culture dishes. Ten nymphs per replicate and three replicates per treatment were used except for experiments with aluminum, where two replicates per treatment were used.

Numbers of nymphs remaining alive in each dish and presence or absence of successful molting were recorded every 4 h until 12 h, and every 12 h thereafter. Dead individuals were identified by lack of response to tactile stimulus.

Experiment E

The effect of pH alone was examined with bioassays at pH 3.0, 4.0, 5.0, 6.0, and 7.5 (control).

Experiment F

The effects of aluminum and pH were examined with bioassays at pH 4.0 and 5.5 at Al concentrations of 0 (control), 100, and 500 $\mu\text{g l}^{-1}$.

Analyses of test waters during Experiments C, D, and F are shown in Table 1.

Statistical analysis

All eggs and nymphs, as well as pH, Al, and Ca treatments were randomly assigned to dishes in each experiment. Differences between treatments in egg hatching and larval mortality were assessed using analysis of variance (ANOVA or GLM), and Duncan's multiple range test (DMRC) was used to compare means. One-way ANOVA (GLM) was used in Experiments A, B, and E, two-way ANOVA in Experiments B and C and three-way ANOVA in Experiment D. The arcsine root transformation was used to normalize variance when data were in the form of percentages. The 96 h LC50 of pH for nymphs was calculated using the POLO program (Russell *et al.*, 1977).

Results

Effects of pH and duration of exposure on eggs (Experiments A and B)

Immediate exposure of eggs to decreasing pH resulted in decreasing ($p < 0.01$) percent successful hatch and increasing ($p < 0.01$) percent partial hatch (Table 2). At pH 3.0 many eggs that did not develop deteriorated rapidly and were floating at the water surface within 24 h; many of the surviv-

Table 2. Mean percent successful hatch \pm SE, partial hatch \pm SE and days to first successful hatch \pm SE of *C. triangulifer* eggs exposed to pH 3.0, 4.0, 5.0, 6.0 and 7.5 (control). Eggs were either immediately exposed after oviposition or exposure was delayed for 7 d while eggs were held at pH 7.5.

pH	% successful hatch		% partial hatch		Days to 1st hatch
	Imm. exp.	Del. exp.	Imm. exp.	Del. exp.	Imm. exp.
3.0	0 \pm 0a*	0 \pm 0a	42.3 \pm 6.6a	69.0 \pm 4.4a	No successful hatch
4.0	4.9 \pm 2.8b	18.3 \pm 12.0b	49.7 \pm 8.3a	34.7 \pm 14.9b	8.0 \pm 0a
5.0	31.2 \pm 7.7c	45.0 \pm 9.0c	40.3 \pm 12.5b	0 \pm 0c	8.0 \pm 0a
6.0	30.1 \pm 7.2c	43.0 \pm 3.5c	5.3 \pm 1.2c	0 \pm 0c	9.0 \pm 1.0a
7.5	42.0 \pm 5.2c	68.0 \pm 2.9d	0.7 \pm 0.3c	0 \pm 0c	8.3 \pm 0.3a

* Means within a column followed by the same letter are not significantly different ($p > 0.05$, DMRC).

ing eggs failed to develop attachment fibers. At pH 4.0, early embryonic development appeared normal but many eggs failed to develop attachment fibers. Days to first successful hatch did not vary among pH levels (Table 2).

Eggs with delayed exposure to decreasing pH also showed decreasing ($p < 0.01$) percent successful hatch and increasing ($p < 0.05$) percent partial hatch (Table 2). Decrease in time of exposure increased ($p < 0.01$) percent successful hatch and decreased ($p < 0.05$) percent partial hatch.




Fig. 1. Mean percent successful hatch of *C. triangulifer* eggs exposed to acidic waters with calcium and aluminum.

Table 3. Mean percent successful hatch, percent partial hatch, and days to first hatch of *C. triangulifer* eggs exposed to pH 4.0 and 5.5 with Ca and Al. Mean values \pm SE for each factor, pooling data across remaining factors.

(1) pH	% successful hatch	% partial hatch	Days to 1st hatch
4.0	17.8 \pm 2.6a*	32.5 \pm 2.2a	11.5 \pm 1.0a
5.5	42.4 \pm 4.1b	13.1 \pm 2.3b	10.4 \pm 0.8a
(2) Al ($\mu\text{g l}^{-1}$)			
0	40.2 \pm 5.5a	20.7 \pm 4.4a	8.3 \pm 0.3a
100	27.3 \pm 3.4b	21.5 \pm 2.7ab	12.1 \pm 1.0b
500	22.8 \pm 5.3b	26.3 \pm 3.5b	12.2 \pm 1.4b
(3) Ca (mg l^{-1})			
0	17.6 \pm 3.9a	15.9 \pm 2.6a	8.6 \pm 0.2a
10	30.2 \pm 4.8b	29.8 \pm 3.9b	14.9 \pm 1.3b
100	42.1 \pm 4.9c	22.8 \pm 3.4c	9.0 \pm 0.7a

* Means followed by the same letter within each column of (1), (2), or (3) are not significantly different ($p > 0.05$, DMRC).




Fig. 2. Mean percent partial hatch of *C. triangulifer* eggs exposed to acidic waters with calcium and aluminum.

Effects of pH, aluminum and calcium on eggs (Experiments C and D)

Increased aluminum alone in acidic waters (Experiment C) did not affect percent successful or partial hatch, or days to first hatch ($p > 0.05$). In contrast, in Experiment D, increased Al ($p < 0.05$) and decreased pH ($p < 0.001$) resulted




Fig. 3. Mean days to first hatch for *C. triangulifer* eggs exposed to acidic waters with calcium and aluminum.

in decreased percent successful hatch. However, increased Ca showed increased percent successful hatch ($p < 0.001$, Table 3, Fig. 1). Percent partial hatch was increased by increased Al ($p < 0.05$) and Ca ($p < 0.001$) and decreased pH ($p < 0.001$) and an Al \times pH interaction was observed ($p < 0.001$, Table 3, Fig. 2).

Days to first hatch showed increase ($p < 0.001$) with Al, Ca and Al \times Ca interaction (Table 3, Fig. 3). Only the moderate level of Ca




Fig. 4. Cumulative percent mortality \pm SE of *C. triangulifer* nymphs exposed to test waters for 96 h. * indicates significant difference ($p < 0.01$).

(10 mg l⁻¹) increased days to first hatch. Difference in pH did not affect days to first hatch ($p > 0.05$).

Effects of pH on nymphs (Experiment E)

Decreased pH reduced ($p < 0.01$) survival of nymphs (Fig. 4). At pH 3.0, there was 100% mortality after 4 h and at pH 4.0 90% mortality after 36 h. There were no differences in mortality at pH 5.0 to pH 7.5. Surviving nymphs in all treatments molted successfully. The calculated 96 h LC50 was pH 4.74 ($n = 150$, slope = 1.03 ± 0.13 , 95% CL = $4.51 \leq 4.74 \leq 4.94$). Accordingly, subsequent experiments were performed within the pH 4.0–5.5 range.

Effect of pH and aluminum on nymphs (Experiment F)

There was an 8–22% lower mortality of nymphs exposed to acidic waters with Al than of nymphs exposed to acidic waters without Al (Table 4). No statistical analysis was attempted with these data as there were only two replicates per treatment. Surviving nymphs in all treatments molted successfully.

Discussion

Gibbs (1977) reported 86% successful hatching of eggs in pond water from which *C. triangulifer*

was collected. Thus exposure of *C. triangulifer* eggs to such dilute test waters may have reduced successful hatching in control (Table 2) as well as experimental treatments.

Our results consistently indicated that decreasing pH decreased successful hatching and increased partial hatching in *C. triangulifer* (Experiments A, B, C and D). All hatches that occurred at pH 3.0 were partial hatches. There were high levels of partial hatches and low levels of successful hatches at pH 4.0 but between pH 5.0 and 7.5 the percent successful hatch remained consistently high. This suggested that hatching of *C. triangulifer* was impaired below pH 5.0. This impairment may have resulted from the direct effect of acidity on the developing embryo, or the nymph as it attempted to break free of the chorion; acidity may also have acted indirectly by affecting the hatching process.

Evidence for a direct effect of low pH on developing embryos of *C. triangulifer* was the increased successful hatching and decreased partial hatching when time of exposure of eggs to acidic conditions was delayed and short-term rather than immediate and long-term (Experiment B). This suggested a direct effect on the embryo related to time of exposure. If egg membranes protect the developing embryo from acidic conditions, the internal pH of eggs would be expected to be different from that of the surrounding water; this hypothesis has not been tested for mayfly eggs.

High levels of partial hatching occurred at low pH, with nymphs dying within the egg or still attached to the chorion by the caudal filaments. This phenomenon was also observed at pH 4.0 and 4.5 in eggs of *Baetis flavistriga* but not in eggs of *Leptophlebia cupida*, *Habrophlebia vibrans* or *Stenonema femoratum* (Rowe *et al.*, 1988b). In *B. flavistriga*, significant mortality of nymphs occurred before the hatch was complete at pH 4.0 (91%) and 4.5 (12%). We believe that the 'incomplete hatch' of Rowe *et al.* (1988b) is equivalent to the 'partial hatch' of this study. Partial hatching has also been reported in fish eggs, with 69% of salmon and 17% of sea-run brown trout dying with the head still attached to chorion at pH 4.1 (Johansson *et al.*, 1977). This hatching failure in

Table 4. Mean ($n = 2$) percent mortality of *C. triangulifer* nymphs exposed to pH 4.0 and 5.0 at aluminum concentrations of 0, 100 and 500 $\mu\text{g l}^{-1}$ for 96 h.

Al ($\mu\text{g l}^{-1}$)	pH	
	4.0	5.0
0	28.6	15.5
100	8.3	5.6
500	6.2	7.1

fish eggs may have resulted from the inactivation of a choriolytic enzyme at reduced pH (Runn *et al.*, 1977; Peterson *et al.*, 1980). Hatching enzymes have been identified in insects (Hinton, 1981) but their presence or function in *C. triangulifer* eggs is unknown.

Reduced successful hatching can also be related to disruption of the mechanical aspects of hatching. In amphibians (Gosner & Black, 1957; Pough, 1976; Dunson & Connell, 1982) the perivitelline space is reduced at low pH leaving little room for muscular movements of the embryo; hatching failure or partial hatching may occur.

Our evidence of possible mechanical disruption of the hatching process was limited to the observation that the egg attachment fibers appeared less abundant at pH 3.0 and 4.0 than at pH 5.0–7.5. This may have contributed to partial hatching by reducing the stability of the substrate from which the nymphs must extricate themselves during hatching.

When levels of both Ca and Al were varied in acidic waters, increased concentrations of Al appeared detrimental, as successful hatching was decreased and partial hatching increased (Experiment D). Also, there was an Al \times pH interaction which resulted in increased partial hatch at the higher pH (5.5) level. Elevated Ca, however, increased both successful and partial hatches. Calcium additions may have had an ameliorating affect on H⁺ stress on *C. triangulifer* eggs by decreasing membrane permeability, which could decrease the rate of internal ion loss. Calcium has been shown to modify the effect of Al at low pH in fish. Brown (1983) reported no or low survival of brown trout fry with 0.25 mg l⁻¹ Ca and almost complete survival with 2 mg l⁻¹ Ca after 16 d at pH's of 4.5–5.4 with 250 μ g l⁻¹ Al. Mortality data for *Baetis rhodani* indicate that nymphs collected from high Ca sites were less sensitive to acidic water with Al than those from low Ca sites (McCahon & Pascoe, 1989).

In contrast to the results with pH alone (Experiment A), moderate levels of Ca (10 mg l⁻¹) increased number of days to first hatch; in addition, days to first hatch was higher at 500 μ g l⁻¹ than at 100 μ g l⁻¹ Al (Experi-

ment D). We have no explanation for this. These experiments were conducted using eggs from overwintering nymphs and Sweeney & Vannote (1984) suggest that eggs from this generation hatch with less synchrony than eggs produced by the summer generations. It is uncertain if this was the case in this instance.

Whatever the cause, increased number of days to first hatch effectively increased experimental exposure times for *C. triangulifer* eggs. We have demonstrated that longer exposures to acidic waters decreased successful hatch and increased partial hatch (Experiment B). Therefore, detrimental effects of H⁺ elevations were exacerbated by increased days to first hatch. This phenomenon, combined with the lack of any Al effects observed in Experiment C, leads us to conclude that Al has little or no independent toxicity to *C. triangulifer* eggs.

Small nymphs showed 100% mortality after 4 h at pH 3.0 and 90% mortality after 36 h at pH 4.0 (Experiment E). This susceptibility may have contributed to the low rate of successful hatch and the high rate of partial hatch at pH 3.0 and 4.0 by directly affecting the nymphs after the chorion was ruptured.

The 96 h LC50 of pH 4.74 indicated that *C. triangulifer* was slightly less tolerant of low pH than *Ephemerella subvaria*, which Bell & Nebeker (1969) found to be the least tolerant mayfly (LC50 = pH 4.65) in their experiments. Tartar *et al.* (1982) found three species of *Baetisca* to be more tolerant than *C. triangulifer* with 96-h LC50's of pH 3.6 for *B. lacustris*, pH 3.5 for *B. carolina* and pH 3.2 for *B. berneri*. Mortalities of *C. triangulifer* nymphs were comparable to those observed by U. Matthias (cited in Okland & Okland, 1986) for *Baetis* spp. nymphs at similar pH's.

Disruption of ionic regulation in acid-stressed mayflies has been indicated by a loss of whole-body sodium and chloride and a correlated mortality (Rowe *et al.*, 1988a, and 1989). Our observations that acid stressed nymphs continued to molt successfully agree with those of Rowe *et al.* (1988a) who reported that molting was more frequent in low pH treatments and was associated

with mortality and large losses of sodium and chloride ions. Rowe *et al.* (1988a, and 1989) suggest that molting is probably a mechanism to inhibit ion loss and not a cause of mortality and may be associated with the increase in number of chloride cells during ionic stress.

Small nymphs such as we used in our experiments are reported to be more sensitive to low pH than larger nymphs (Allard & Moreau 1987). However, Rowe *et al.* (1989) found that nymphs within the size range of 2–14 mg dry wt showed no size related ion losses at pH 3.5.

Aluminum at 100 and 500 $\mu\text{g l}^{-1}$ appeared to ameliorate the effect of pH on mortality of nymphs of *C. triangulifer* at pH 4.0 and 5.0 (Experiment F). Although Al toxicity to fish has frequently been reported at low pH (see review by McCahon *et al.*, 1987) ameliorating effects of Al on H^+ stress in brook trout fry (*Salvelinus fontinalis*) at pH 4.0 have also been reported by Schofield (1977) and in brown trout by Muniz & Leivestad (1980). Insects, including *C. triangulifer*, appear to be less sensitive than fish to Al toxicity. Increased gill mucus production, which has been observed in fish, was not found in mayfly nymphs (*Baetis rhodani*, *Ecdyonurus venosus*) exposed to an Al concentration of 350 $\mu\text{g l}^{-1}$ at pH 5.0 (McCahon *et al.*, 1987). Cook & Haney (1984) found that of the nine species tested (five caddisfly species, two mayfly species, a stonefly, and a beetle) Al additions caused increased mortality only in the stonefly *Nemoura nigratta* and the caddisflies of the genus *Macronema*.

Evidence of significant reduction of successful hatching and survival of nymphs below pH 5.0 suggested that *C. triangulifer* is one of the more sensitive mayflies for which data are available. Other mayflies showing comparable sensitivity, *Baetis lapponicus*, *B. macani* (Okland & Okland, 1986) and *B. flavistriga* (Rowe *et al.*, 1988b), are also members of the family Baetidae, indicating that this taxon may be especially vulnerable to habitat acidification. Our observations that *C. triangulifer* eggs and small nymphs are tolerant of short-term exposures to levels of aluminum occurring in Maine surface waters suggest that this species could survive episodic Al increases

such as those following snowmelt in streams. However, extended exposures of previously acid-stressed eggs could cause a detrimental Al \times pH interaction, resulting in *C. triangulifer* recruitment reduction or failure. These negative effects are most pronounced in low ionic-strength water and can be ameliorated to a certain extent by the addition of calcium.

Acknowledgements

We thank Connie Stubbs, Brian Molloy, Mike Schorr, and Terry Mingo for assistance in the laboratory, and Susan Elias, Myron Buck, and Scott Denning for water chemistry analysis. Terry Haines, Steve Kahl, Randy Alford, Ellie Groden and Bill Halteman offered comments on an earlier version of the manuscript. This work was funded in part by the Department of the Interior, U.S. Geological Survey, through the Maine Land and Water Resources Center and the Maine Agricultural Experiment Station, University of Maine.

References

- Allard, M. & G. Moreau, 1987. Effects of experimental acidification on a lotic macroinvertebrate community. *Hydrobiologia* 144: 37–49.
- Baker, J. P. & C. L. Schofield, 1982. Aluminum toxicity to fish in acidic waters. *Wat. Air Soil Pollut.* 18: 289–309.
- Bell, H. L. & A. V. Nebeker, 1969. Preliminary studies on the tolerance of aquatic insects to low pH. *J. Kan. Entomol. Soc.* 42: 230–236.
- Brown, D. J. A., 1983. Effect of calcium and aluminum concentrations on the survival of brown trout (*Salmo trutta*) at low pH. *Bull. envir. Contam. Toxicol.* 30: 582–587.
- Cook, W. & J. Haney, 1984. The acute effects of aluminum and acidity upon nine stream insects. *Water Resource Research Center, Durham, NH, Res. Rep.* 46, 33 pp.
- Dunson, W. D. & J. Connell, 1982. Specific inhibition of hatching in amphibian embryos by low pH. *J. Herpetol.* 16: 314–316.
- Fiance, S., 1978. Effects of pH on the biology and distribution of *Ephemerella funeralis* (Ephemeroptera). *Oikos* 31: 332–339.
- Gibbs, K. E., 1977. Evidence for obligatory parthenogenesis and its possible effect on the emergence period of *Cloeon triangulifer* (Ephemeroptera: Baetidae). *Can. Entomol.* 109: 337–340.

- Gosner, K. L. & I. H. Black, 1957. The effect of acidity on the development and hatching of New Jersey frogs. *Ecology* 38: 256–262.
- Haines, T. H. & J. J. Akielaszek, 1983. A regional survey of chemistry of headwater lakes and streams in New England: vulnerability to acidification. U.S. Fish and Wildlife Service, Eastern Energy and Land Use Team, FWS/OBS-80/40.15, 141 pp.
- Hinton, H. E., 1981. *Biology of insect eggs*. Pergamon Press, Oxford, 1125 pp.
- Johansson, N., P. Runn & G. Milbrink, 1977. Early development of three salmonid species in acidified water. *Zoon* 5: 127–132.
- Keller, W., 1983. Spring pH and alkalinity depressions in Lake Superior tributaries. *J. Great Lakes Res.* 9: 425–429.
- McCahon, C. P. & D. Pascoe, 1989. Short-term experimental acidification of a Welsh stream: toxicity of different forms of aluminum at low pH to fish and invertebrates. *Arch. envir. Contam. Toxicol.* 18: 233–242.
- McCahon, C. P., D. Pascoe & C. McKavanagh, 1987. Histochemical observations on the salmonids *Salmo salar* L. and *Salmo trutta* L. and the ephemeropterans *Baetis rhodani* (Pict.) and *Ecdyonurus venosus* (Fabr.) following a simulated episode of acidity in an upland stream. *Hydrobiologia* 153: 3–12.
- Muniz, I. P. & H. Leivestad, 1980. Toxic effects of aluminum on the brown trout, *Salmo trutta* L., p. 320–321. In D. Drablos & A. Tollan (eds.), *Ecological Impact of Acid Precipitation*. SNSF Project, Sandfjord, Norway.
- Okland, J. & K. A. Okland, 1986. The effects of acid deposition on benthic animals in lakes and streams. *Experientia* 42: 471–486.
- Ormerod, S. J., P. Boole, C. P. McCahon, N. S. Weatherley, D. Pascoe & R. W. Edwards, 1987. Short-term experimental acidification of a Welsh stream: Comparing the biological effects of hydrogen ions and aluminum. *Freshwat. Biol.* 17: 341–356.
- Peterson, R. H., P. G. Daye & J. L. Metcalfe, 1980. Inhibition of Atlantic salmon (*Salmo salar*) hatching at low pH. *Can. J. Fish. aquat. Sci.* 37: 770–774.
- Pough, F. H., 1976. Acid precipitation and embryonic mortality of spotted salamanders, *Ambystoma maculatum*. *Science* 192: 68–70.
- Rosseland, B. O. & O. K. Skogheim, 1982. Physiological stress and mortality of Atlantic salmon, *Salmo salar* L., in acid water with high levels of aluminum. International Council for the Exploration of the Sea, Anadromous and Catadromus Fish Committee. C.M. 1982/M: 29. 17 pp.
- Rowe, L., M. Berrill & L. Hollett, 1988a. The influence of season and pH on mortality, molting and whole-body ion concentrations in nymphs of the mayfly *Stenonema femoratum*. *Comp. Biochem. Physiol.* 90A: 405–408.
- Rowe, L., J. Hudson & M. Berrill, 1988b. Hatching success of mayfly eggs at low pH. *Can. J. Fish. aquat. Sci.* 45: 1649–1652.
- Rowe, L., M. Berrill, L. Hollett & R. J. Hall, 1989. The effects of short-term laboratory pH depressions on molting, mortality and major ion concentrations in the mayflies *Stenonema femoratum* and *Leptophlebia cupida*. *Hydrobiologia* 184: 89–97.
- Runn, P., N. Johansson & G. Milbrink, 1977. Some effects of low pH on hatchability of eggs of perch, *Perca fluviatilis* L. *Zoon* 5: 115–125.
- Russell, R. M., J. L. Robertson & N. E. Savin, 1977. POLO: A new computer program for probit analysis. *Bull. entomol. Soc. Am.* 23: 209–213
- Schofield, C. L., 1977. Acid snow-melt effects on water quality and fish survival in the Adirondack Mountains of New York State. *Res. Proj. Tech. Compl. Rep. A-072-NY*. Office of Water Research and Technology, Dept. of the Interior, Washington, D.C., 27 pp.
- Stumm, W. & J. J. Morgan, 1981. *Aquatic chemistry*, second ed. Wiley Interscience, New York, NY., 780 pp.
- Sweeney, B. W. & R. L. Vannote, 1984. Influence of food quality and temperature on life history characteristics of the parthenogenetic mayfly *Cloeon triangulifer*. *Freshwat. Biol.* 14: 621–630.
- Tartar, D. C., D. L. Chaffee & K. B. Benson, 1982. Low pH tolerance under laboratory condition of the nymphs of three congeneric mayflies, *Baetisca berreri*, *B. carolina* and *B. lacustris*. *Proc. W. VA. Acad. Sci.* 54(1): 11–18.
- Wiederholm, T., 1984. Responses of aquatic insects to environmental pollution, p. 530–535. In V. H. Resh & D. M. Rosenberg (eds.), *The Ecology of Aquatic Insects*. Praeger Publishers, New York.